	Potential years o	f life lost from cau	uses considered a	menable to healthca	re – adults				Potential years of life lost from causes amenable to healthcare CYP
OUTCOME MEASURES SYSTEM METRICS	CHD Under 75s Mortality Rate from all cardio vascular disease Under 75 mortality rate from cardiovascular disease considered preventable	Stroke (Preventable mortality)	Diabetes (Preventable mortality)	Cancer Under 75s mortality rate from cancer Under 75 mortality rate from cancer considered preventable from Cancer 1 & 5 year survival from cancers	Chronic Respiratory Under 75s Mortality Rate from respiratory disease Under 75 mortality rate from respiratory disease considered preventable	CKD (Preventable mortality)	Mental Health Premature mortality in adults with severe mental illness Excess mortality rate in adults with severe mental illness Suicide rate Deaths from drug misuse	MSK Hip fractures in people aged 65 and over	Maternal & Child Health Infant mortality rate Good Level of Development age 2.5 and 5 years
URGENT CARE Emergency admissions for ambulatory care and urgent care sensitive conditions Avoidable Hospital admissions Readmissions	Emergency MI admissions	Emergency Admissions due to Stroke		% diagnosed on A& E attendance			Mental Health Streaming A&E attendances/Emergency hospital admissions for self-harm	Emergency admission due to falls	Unplanned hospitalisation for asthma, diabetes and epilepsy in under 19s Accidents & unintentional injury Emergency admissions for children with lower respiratory tract conditions
PLANNED CARE Unmet need for health care Stage of presentation, RTT waits, DNAs,)	Diagnostic waits			Cancers diagnosed at early stage Treatment within 62 days			IAPT access IAPT recovery EIP Waiting Times Inpatient admissions Out of area placements Admission episodes for alcohol related conditions		Waits LAC, CIN, CSC Tooth extractions due to decay for children admitted as inpatients to hospital, aged 10 years and under Breast feeding initiation

Independent Living among those with Multi-morbidity  Health related quality of life for people with Long term conditions  Proportion of people feeling supported to manage their condition  Employment of People with Long term conditions  Health related quality of life for carers  Proportion of older people still home 91 days post discharge			Diabetes patients achieving NICE recommended targets				Employment of people with mental illness Successful completion drug treatment Successful completion alcohol treatment	Rapid response – response to falls/admission avoidance Hip fracture: proportion of people recovering to previous levels of mobility/walking ability at 30 day and 120 days	Social Prescribing Referrals % of children receiving 6-8 week review Breast feeding at 6-8 weeks New birth visits within 14 days
PRIMARY CARE Proportion of people feeling supported to manage their condition	NHS Health Checks Expected vs observed CVD registers % treated to target Referrals to Live Well Services 'Social prescribing' referrals and outcomes	Expected vs Recorded Prevalence (Missing Thousands) % achieving best care indicators Referrals to Live Well Services 'Social prescribing' referrals and outcomes	Expected vs Recorded Prevalence (Missing Thousands) % achieving best care indicators Referrals to Live well Services Preventable sight loss — diabetic disease	Screening uptake Two week waits Referrals to Live well Services 'Social prescribing' referrals and outcomes	Expected vs Recorded Prevalence (Missing Thousands) % achieving best care indicators Referrals to Live Well Services 'Social prescribing' referrals and outcomes	Expected vs Recorded Prevalence (Missing Thousands) % achieving best care indicators Referrals to Live Well Services 'Social prescribing' referrals and outcomes	Dementia diagnosis rate Dementia care plan reviewed within 12 months People with SMI received health check Referrals to Live Well Services 'Social prescribing' referrals and outcomes	Osteoporosis Screening and management Referrals to Live Well Service 'Social prescribing' referrals and outcomes	Childhood Imms LARC Tooth extractions due to decay for children admitted to hospital Child development – communication skills at 2.5 years Personal social skills at 2.5 years

POP HEALTH Smoking prevalence % classed as Overweight or obese % Physical Activity levels % Flu uptake	% achieving health goals with Live Well Service	% achieving health goals with Live Well Service	% achieving health goals with Live Well Service	% achieving health goals with Live Well Service	% achieving health goals with Live Well Service	% achieving health goals with Live Well Service	% achieving health goals with Live Well Service Self reported wellbeing	% achieving health goals with Live Well Service % injuries form falls in people aged 65+	Number of referrals into lifestyle services % children aged reception & year 6 classified as overweight or obese % physically active children Smoking prevalence at age 15 Maternal smoking at time of delivery Maternal obesity Alcohol at booking
									,